



Functionalized Vinyl Boronates for C-C couplings

Introduction


The Suzuki–Miyaura cross-coupling of boronic acid derivatives is one of the most used reactions in organic and medicinal chemist's toolbox. The rapid advancement of this method resulted in its efficient application for the late-stage modification of biologically active substrates and construction of combinatorial libraries. One of the recent trends in the field of organoboron reagents is related to the shift from aromatic compounds towards cyclic sp^3 -enriched structures, which comply with criteria of lead-oriented synthesis.¹⁻⁶ In this context, *Enamine* offers a library of alkenylboronic esters for metal-mediated couplings.



Case studies



We offer: more than 50 of vinyl boronates from stock on a 5-10 g scale.



References

1. M. Kovalenko et al. *Eur. J. Org. Chem.* **2019**, 5624.
2. D. G. Brown et al. *J. Med. Chem.* **2016**, 59, 4443.
3. S. D. Roughley et al. *J. Med. Chem.* **2011**, 54, 3451.
4. P. Schäfer et al. *Nat. Commun.* **2018**, 9, 16216.
5. P. S. Campbell et al. *Chem. Commun.* **2018**, 54, 46.
6. A. Nadin et al. *Angew. Chem. Int. Ed.* **2012**, 51, 1114.



Search & Buy on-line at EnamineStore.com
Look for more at Chem-Space.com

BB@enamine.net, www.enamine.net