


# Regioselectivity study of Beckman reaction as a pathway for synthesis of bicyclic anilines and benzylamines




Bohdan Yu. Solod, Yevhen M. Ostapchuk, Mykhailo V. Vovk, Tetiana V. Druzhenko, Dmytro M. Volochnyuk, Serhiy V. Ryabukhin

## Background of the project

The Beckmann rearrangement is a highly versatile and industrially important reaction that plays a crucial role in the production of polymers & pharmaceuticals due to its ability to efficiently produce lactams and amides




**OBSTACLE:** Low regioselectivity of the Beckmann rearrangement of bicyclic aryl ketones poses a great problem to its wide application in synthetic organic and process chemistry:



## THIS PROJECT:


- bicyclic aryl ketones with cyclopentane and cyclohexane moieties
- A wide range of EWG, 'neutral', and EDG substituents placed in the aromatic ring in different positions
- thoroughly optimized rearrangement conditions



## Contact

Sergey V. Ryabukhin, Prof. Dr. Sci. [s.v.ryabukhin@gmail.com](mailto:s.v.ryabukhin@gmail.com); Dmytro M. Volochnyuk, Prof. Dr. Sci. [d.volochnyuk@gmail.com](mailto:d.volochnyuk@gmail.com)

## Optimization of the rearrangement conditions and 'structure/direction' relationships



- Multiparameter optimization of rearrangement conditions includes different reagents, solvents, temperature modes, and reaction times
- Identified the three reaction conditions providing the best results, which were selected for the further substrates screening

product 1	product 2	product 1	product 2	product 1	product 2	product 1	product 2	product 1	product 2	product 1	product 2	product 1	product 2
33%	67%	0%	100%	12%	88%	25%	75%	100%	0%	100%	0%	100%	0%
18%	82%	16%	84%	0%	100%	73%	27%	100%	0%	100%	0%	100%	0%
7%	93%	0%	100%	0%	100%	81%	19%	100%	0%	100%	0%	100%	0%
STARTING MATERIAL		60%	40%	0%	100%	56%	44%	100%	0%	100%	0%	100%	0%
DECOMPOSITION PRODUCTS		17%	83%	0%	100%			100%	0%	100%	0%	100%	0%

INVERSE DIRECTION OF THE REARRANGEMENT

## Future horizons

- DFT calculations and building a predicting model for the studied aryl ketones allowing for the efficient expansion of the lactams chemical space
- To involve in the project a wide range of heteroaromatic ketones (208 substrates in total,  $n = 1, 2$ , Q = heteroatom, A = heteroatom or carbon)

